

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**

12

EUROPEAN PATENT APPLICATION

21 Application number: 83305042.6

22 Date of filing: 01.09.83

51 Int. Cl.<sup>3</sup>: C 12 N 5/00  
C 12 P 1/00, C 12 Q 1/00  
//A61K39/395

30 Priority: 01.09.82 US 413861

43 Date of publication of application:  
28.03.84 Bulletin 84/13

64 Designated Contracting States:  
DE GB

71 Applicant: Sloan-Kettering Institute For Cancer  
Research  
1275 York Avenue  
New York New York 10021(US)

72 Inventor: Calmcross, Gregory J.  
229 Sherwood Avenue  
London Ontario(CA)

72 Inventor: Mattes, Jules M.  
178/10 Wexford Terrace  
Jamaica Estates New York 11432(US)

72 Inventor: Beresford, Richard H.  
Bay Avenue  
Centre Island New York 11771(US)

72 Inventor: Albino, Anthony P.  
504 East 81st Street  
New York New York 10028(US)

72 Inventor: Houghton, Alan N.  
450 East 63 Street  
New York New York 10021(US)

72 Inventor: Lloyd, Kenneth O.  
4525 Henry Hudson Parkway  
Bronx New York(US)

72 Inventor: Old, Lloyd J.  
600 West End Avenue  
New York New York(US)

74 Representative: Bernard, Alan Peter et al,  
F.J. CLEVELAND & COMPANY 40/43 Chancery Lane  
London WC2A 1JQ(GB)

54 Cell surface antigens of human astrocytoma.

57 Method of forming an antibody producing hybridoma cell line by fusing a myeloma derived cell strain with splenocytes derived from BALB/c mice immunized with human astrocytoma tumor cells, the hybridoma cell line formed, and the monoclonal antibodies generated by said cell line. A method of phenotyping astrocytoma tumor cells comprising determining the reaction of said cells to various phenotyped monoclonal antibodies to astrocytoma tumor cells is also provided.

EP 0 104 014 A2

- 1 -  
DESCRIPTION

The present invention is concerned with various surface antigens of human cancer identified by mouse monoclonal antibodies and more particularly, with mouse monoclonal antibodies recognizing various antigens expressed by human malignant astrocytoma.

Astrocytoma tumors are normally found only in the brain and are difficult to diagnose in a manner wherein a useful prognosis can be generated and the most effective treatment predicted. The present invention provides monoclonal antibodies which may be used in identifying or phenotyping astrocytoma tumors whereby the course of the disease can be predicted and suitable therapy, e.g. chemotherapy versus radiotherapy -- can be accomplished.

Furthermore, the present invention provides monoclonal antibodies which may be coupled with various side or toxic agents which then may be used for actual treatment or containment of astrocytoma tumors. Imaging of the tumor may also be possible using various radio labels; however, other means which are believed to be more practical, are available for this purpose.

The present invention also provides a process by which the inventive materials can be raised by sensitization of mice to raise the antibodies and the application of hybridoma fusion techniques to immortalize the cell line.

- 2 -

Work has been done in general with cell surface antigens of human cancers and cell surface antigens of human malignant melanoma and human renal cancer identified by mouse monoclonal antibodies have been described in the literature (1, 2).

Abbreviations used in this disclosure: GFA, glial fibrillary acidic protein; Ab, monoclonal antibody; EBV, Epstein-Barr virus; NP-40, Nonidet P40; conA, concanavalin A; PMSF, Phenylmethylsulfonylfluoride; PAGE, polyacrylamide gell electrophoresis; 2-D, two-dimensional electrophoresis; IEF, isoelectric focusing.

In the following antibodies in accordance with the invention are referenced by codes, e.g. Ab A010. Hybridomas producing these antibodies have been deposited at the American Type Culture Collection, and the following table lists the deposition Number of the hybridoma and the corresponding antibody reference.

Antibody Reference	ATCC Deposition No.
AJ 2	HB 8338
AJ 8	HB 8339
AJ 9	HB 8340
AJ 10	HB 8341
AJ 17	HB 8342
AJ 60	HB 8343
AJ 225	HB 8344
AJ 425	HB 8345
AO 10	HB 8346
AO 50	HB 8347
AO 92	HB 8348
AO 122	HB 8349

The surface antigens of cultured human malignant astrocytomas were analyzed using mouse monoclonal antibodies. BALB/c mice were immunized repeatedly with either SK-MG-1 [a glial fibrillary acidic protein (GFA)-negative astrocytoma line] or SK-A02 [a GFA-positive astrocytoma line]. Following fusion with NS/1 mouse myeloma cells, 12 antibody-producing clones were selected for detailed study. Serological analysis permitted the identification of 9 distinct antigenic systems. Four antibodies (Ab AJ225, Ab A010, Ab AJ8, Ab A0122) identified cell surface antigens preferentially expressed on tumors of neuroectodermal origin, and these antibodies subdivided the astrocytoma panel into distinguishable subsets. The determinants detected by Ab A010 and Ab AJ8 showed mutually exclusive expression on the astrocytoma lines. The A010 and AJ8 phenotypes appeared to reflect the differentiation state of the cultured cells; 4/7 A010-positive astrocytomas expressed GFA, an intracellular astrocyte differentiation antigen, whereas all AJ8-positive astrocytomas (9/9) were GFA-negative. Five antibodies (Ab AJ10, Ab AJ9, Ab AJ17, Ab AJ425, Ab AJ2) recognized determinants widely distributed on normal and malignant cells. AJ2 was highly immunogenic in mice, and antibody to AJ2 was commonly found in the serum of mice immunized with human cells. Four antibodies defined in this study precipitated proteins from reduced preparations of radioisotope-labelled SK-MG-1 and SK-A02 cells: Ab AJ225 ( $M_r$  145,000), Ab A0122 ( $M_r$  265,000), Ab AJ10 ( $M_r$ s 195,000, 165,000) and Ab AJ2 ( $M_r$ s 170,000, 140,000, 140,000, 28,000).

### Tissue Culture

Astrocytoma and other human tumor cell lines and short-term cultures of normal human skin fibroblasts and kidney epithelial cells have been described (3,4,5).

### Serological Procedures

Direct serological tests were performed using an anti-mouse Ig mixed hemadsorption assay (1). Direct test conditions and absorption procedures were identical to those previously described for the protein A-mixed hemadsorption assay (4). Heat stability of the antigenic determinants was assessed by heating the cell suspension to 100°C for 5 minutes and then testing for residual antigenic activity in absorption tests. Glial fibrillary acidic protein (GFA) was demonstrated by indirect immunofluorescence using monospecific rabbit antiserum provided by Dr. L.F. Eng.

### Immunizations

BALB/c mice were immunized with either the GFA-negative astrocytoma line SK-MG-1 [designated AJ in a previous publication (3)] or the GFA-positive astrocytoma line SK-A02 (established by J.R. Shapiro and W.R. Shapiro, Laboratory of Neuro-Oncology, Sloan-Kettering Institute). For the initial immunization,  $1 \times 10^7$  astrocytoma cells were injected subcutaneously with Freund's complete adjuvant. Five to ten subsequent immunizations were carried out at 2-week intervals by intraperitoneal inoculation of  $1 \times 10^7$  tumor cells in the absence of adjuvant. Immunized mice were sacrificed 3 days after the last injection.

Production of Mouse Monoclonal Antibodies

The fusion of immune spleen cells with mouse myeloma MOPC-21 NS/1 cells (ratio 5:1) was performed as described (1,2). Fused cells were grown in selective medium and subcloned by limiting dilution as previously described (1,2). For initial screening, supernatants were tested for antibody activity on a panel of cultured cells consisting of 3 astrocytoma cell lines (including the immunizing line), 2 melanomas, 5 carcinomas and adult and fetal skin fibroblasts. Antibody subclass was determined by double diffusion in agar with anti-Ig heavy-chain-specific reagents (Bionetics, Kensington, MD). Cultures of cloned hybridomas were injected subcutaneously into nu/nu mice (NIH Swiss Background). Sera from mice with progressively growing tumors were used for serological and biochemical characterization.

Immunoprecipitation Procedures

Antibodies were tested for precipitating activity using radiolabelled antigen from detergent solubilized extracts of the immunizing cell line. Three different labelling procedures were used. Labelling with



[<sup>3</sup>H]glucosamine (New England Nuclear, 30-60 Ci/mmole) or with [<sup>35</sup>S]methionine (New England Nuclear, 1000 Ci/mmole) and extraction with Nonidet P40 (NP-40) buffer were carried out as described previously (1,2). In some experiments the [<sup>35</sup>S]methionine-labelled extract was fractionated on a 1 ml concanavalin A (conA)-Sepharose (Pharmacia, Piscataway, NJ) column, using 0.15 M NaCl, 0.01 M Tris HCl pH 7.3, 0.1% NP-40 as column buffer, and eluting with 0.2 M  $\alpha$ -methyl D-mannoside. <sup>125</sup>I-labelling of solubilized membrane preparations followed Brown et al (6), except that the gel filtration step before iodination was omitted. Membrane preparation was carried out according to Natori et al (7), except that the buffer during disruption was supplemented with 10 mM MgCl<sub>2</sub> and 2 mM phenylmethylsulfonylfluoride (PMSF). Protein-conjugated <sup>125</sup>I was estimated by counting samples precipitated with cold 10% (w/v) trichloroacetic acid and then washed with ethanol and acetone.

Radioimmunoprecipitation procedures with [<sup>3</sup>H]glucosamine- and [<sup>35</sup>S]methionine-labelled samples were carried out as described (1,2). For <sup>125</sup>I-labelled samples, aliquots (200  $\mu$ l) were first precleared of non-specific binding material by treating with nu/nu mouse serum (1  $\mu$ l), rabbit anti-mouse IgG (15  $\mu$ l) (Cappel Laboratories, Cochranville, Pa) and Staphylococcus aureus (15  $\mu$ l) (Bethesda Research Laboratories, Bethesda, Md). To aliquots (200  $\mu$ l) of the precleared supernatant solution ( $5 \times 10^5$  cpm <sup>125</sup>I-protein), 0.1 or 1  $\mu$ l antibody and 15  $\mu$ l rabbit anti-mouse IgG were added. Immune complexes were isolated with S. aureus as described (6), except that 0.1 ml of normal rabbit serum was added to the second wash buffer to reduce background binding. Labelled antigen was eluted from the pellet and analyzed by NaDodSO<sub>4</sub>/polyacrylamide gel electrophoresis (PAGE).

- 7 -

and two-dimensional (2-D) electrophoresis as described previously (8,9), except that iodoacetamide (14 mg/ml) was added to the sample buffer when non-reduced samples were analyzed.

## RESULTS

From 4 fusions of NS/1 myeloma with spleen cells from mice immunized with SK-MG-1 (3 fusions) or SK-A02 (1 fusion), 12 antibody-producing clones were selected for detailed analysis (Table 1). The serological specificity of these antibodies was tested on a panel of 49 established human cell lines (Table 2). The antibodies were also tested on short-term cultures of human adult and fetal skin fibroblasts, kidney epithelial cells and melanocytes. In most cases, serological analysis consisted of both direct and absorption tests. Melanocytes were studied only by direct test, and lymphoblastoid lines, erythrocytes, adult brain and fetal brain were analyzed only by absorption tests.

Monoclonal antibodies Ab AJ225, Ab A010, Ab AJ8, Ab A0122, Ab AJ10, Ab AJ9, Ab AJ17, Ab AJ425 and Ab AJ2 defined 9 distinct cell surface antigenic systems. They have been selected for detailed presentation (Table 2). [Of the remaining 3 monoclonal antibodies listed in Table 1, Ab A050 and Ab A092 were serologically related to Ab A0122, and Ab AJ60 was similar to Ab AJ10].

### AJ225 Antigenic System

Direct tests and absorption analysis with Ab AJ225 indicated that the determinant detected by this antibody was largely restricted to astrocyte cell lines (Table 2). Although all astrocytoma lines absorbed Ab AJ225

reactivity, the titers in direct serological tests permitted a division of cultured astrocytomas into two groups based upon quantitative differences in antigen expression; 12/16 tumors were high expressors and 4/16 low expressors. The only other tumor lines expressing high levels of AJ225 were 1/10 melanomas and 1/17 epithelial cancers; 2/4 renal carcinomas expressed low levels of the antigen demonstrable by absorption tests. In addition, a T-cell leukemia (MOLT-4) absorbed Ab AJ225 reactivity. EBV-transformed B-cells, adult and fetal skin fibroblasts, kidney epithelial cells, and homogenates of adult and fetal brain did not absorb. The results of direct serological tests on melanocytes suggest low levels of AJ225 expression on this normal cell type.

Ab AJ225 identified a heat-labile determinant and immunoprecipitated a protein with  $M_r$  145,000 from  $^{125}\text{I}$ -labelled SK-MG-1 (Fig. 1). This band was not detected in precipitates from cells labelled with  $[^3\text{H}]$ glucosamine or  $[^{35}\text{S}]$ methionine. In some experiments, this antigen appeared as a closely spaced doublet. The  $pI$  of the 145,000  $M_r$  component was 4.8. Without reduction a single band with  $M_r$  150,000 was identified relative to reduced standards. The inability to precipitate the AJ225 antigen after metabolic labeling raised the possibility that this determinant was a fetal calf serum component adsorbed to the cell surface. However, the highly restricted distribution of this antigen, the failure of fetal calf serum to inhibit Ab AJ225, and the ability of fresh astrocytoma tissue to absorb Ab AJ225 reactivity speak strongly against this possibility.

A010 and AJ8 Antigenic Systems

The A010 and AJ8 antigenic systems are described together because, with two exceptions, they subdivided 16 cultured astrocytomas into mutually exclusive A010-positive and AJ8-positive subsets. The exceptions were SK-MG-10 which expressed neither antigen and SK-MG-12 which expressed both.

The A010 antigen was identified on 7/16 astrocytoma lines. As shown in Table 2, 4/7 A010-positive astrocytomas were GFA-positive. The antigen was also demonstrated on 3/10 melanomas, 2/2 neuroblastomas, and a T-cell leukemia. Direct serological tests failed to demonstrate the A010 determinant on epithelial cancers; however, absorption of Ab A010 reactivity indicated low levels of antigen expression on 2/17 of these lines. EBV-transformed B-cells did not absorb Ab A010 reactivity. The A010 determinant was detected in adult and fetal brain but not identified on adult and fetal skin fibroblasts, kidney epithelial cells or melanocytes.

The AJ8 antigen was detected on 9/16 astrocytoma lines. All AJ8-positive astrocytomas were GFA-negative (Table 2). The AJ8 antigen was also demonstrated on 4/10 melanomas. Direct serological tests failed to demonstrate the AJ8 determinant on epithelial cancers; however, absorption analysis detected low levels of antigen expression on 4/17 cell lines. Neuroblastomas (0/2), EBV-transformed B-cells and a T-cell leukemia did not absorb Ab AJ8 reactivity. AJ8 was detected on adult and fetal skin fibroblasts and melanocytes, but not on cultured kidney epithelial cells or in adult or fetal brain.

The A010 determinant was heat-labile, suggesting that it resided on a protein, but Ab A010 did not precipitate a detectable component from [<sup>3</sup>H]glucosamine-, [<sup>35</sup>S]methionine-, or <sup>125</sup>I-labelled SK-A02 cells. The AJ8

determinant was also heat-labile and could not be precipitated from [ $^3\text{H}$ ]glucosamine-, [ $^{35}\text{S}$ ]methionine-, or  $^{125}\text{I}$ -labelled SK-MG-1 cells.

#### A0122 Antigenic System

The A0122 antigen was found on 9/16 astrocytoma lines (Table 2). The pattern of expression on astrocytomas clearly distinguished the A0122 system from the A0225, A08 or A010 systems. A0122 was strongly represented on 8/10 melanomas. Neuroblastomas (0/2), epithelial cancers (0/17), EBV-transformed B-cells, and a T-cell leukemia did not express A0122. Melanocytes and adult and fetal skin fibroblasts were highly reactive with Ab A0122, and homogenates of adult and fetal brain absorbed Ab A0122 reactivity. A0122 was not detected on kidney epithelial cells or erythrocytes.

Ab A0122 identified a heat-labile determinant and immunoprecipitated a protein complex from  $^{125}\text{I}$ -labelled SK-A02 cells. Four major proteins with  $M_r$ s 265,000, 195,000, 180,000 and 140,000 were identified in reduced preparations. Without reduction, 4 components with  $M_r$ s 255,000, 150,000, 135,000 and 115,000 were seen. Only the 265,000  $M_r$  band was detected after labelling with [ $^{35}\text{S}$ ]methionine. Ab A0122 did not precipitate a detectable component from [ $^3\text{H}$ ]glucosamine labelled SK-A02 cells. Two other monoclonal antibodies, Ab A050 and Ab A092, appeared to recognize the A0122 determinant. The pattern of Ab A050 reactivity was identical to Ab A0122, but Ab A050 did not precipitate any components from  $^{125}\text{I}$ -labelled SK-A02 cells. Ab A092 precipitated the same protein complex as Ab A0122, although minor differences were found in the serological reactivity of the two antibodies.

AJ10 Antigenic System

The AJ10 determinant was found to be widely distributed on normal and malignant cells (Table 2). Despite this broad representation, the AJ10 antigen was not detected on any of the breast or colon cancer cell lines.

Ab AJ10 identified a heat-labile determinant and immunoprecipitated two clearly indentifiable proteins with  $M_r$ s 195,000 and 165,000 from reduced extracts of [ $^{35}$ S]methionine- or  $^{125}$ I-labelled SK-MG-1 cells

Unreduced samples migrated as two bands corresponding to  $M_r$ s 135,000 and 110,000. Ab AJ10 did not precipitate a detectable component from [ $^3$ H]glucosamine-labelled cells. A second monoclonal antibody, Ab AJ60, demonstrated identical serological reactivity to Ab AJ10 but did not precipitate the 195,000/165,000  $M_r$  complex.

AJ2 Antigenic System

AJ2 was demonstrated on all nucleated human cells examined, both normal and malignant (Table 2). Ab AJ2 identified a heat-labile determinant and immunoprecipitated a glycoprotein complex from [ $^3$ H]glucosamine-, [ $^{35}$ S]methionine- and  $^{125}$ I-labelled SK-MG-1 cells. With  $^{125}$ I-labelling and under reduced conditions 3 components were identified by NaDodSO<sub>4</sub>/PAGE:

2 heavy chains with  $M_r$ s 170,000 and 140,000 and one light chain with  $M_r$  28,000

Under non-reduced conditions, the 2 high molecular weight components migrated in a similar fashion, but the light chain was not identified

2-D electrophoresis resolved the 140,000  $M_r$  band into two distinct bands with slightly different molecular weights

The 170,000  $M_r$  component had a pI of 5.2; the two 140,000  $M_r$  chains had isoelectric points of 5.5 and 4.7. The 28,000  $M_r$  light chain focused in

- 12 -

3 major spots at pH 6.2, 5.9 and 5.7.

The light chain was disulfide-linked to only one of the heavy chains. This was demonstrated in two ways. 2-D electrophoresis without reduction resulted in the disappearance of the light chain components and a shift upward of the 140,000  $M_r$  chain with pI 5.5.

Reduction after isoelectric focusing and before NaDodSO<sub>4</sub>/PAGE demonstrated the light chain beneath the 140,000  $M_r$  chain with pI 5.5.

[<sup>35</sup>S]methionine labelled the 4 subunits, but the light chain very weakly. With [<sup>3</sup>H]glucosamine-labelling, only the heavy chains were labelled. The [<sup>35</sup>S]methionine-labelled SK-MG-1 membrane preparation was fractionated on a conA-Sepharose column; Ab AJ2 precipitated the 170,000/140,000/140,000/28,000  $M_r$ s complex from the bound and eluted glycoprotein fraction.

Our experience suggests that AJ2 is highly immunogenic in mice. A range of antisera from mice hyperimmunized with different human cell lines immunoprecipitated a 140,000  $M_r$  component from [<sup>35</sup>S]methionine-labelled cells.

Preclearing with Ab AJ2 removed this  $M_r$ 140,000 component. Preclearing with monoclonal antibodies of the same Ig subclass as Ab AJ2 but directed against other determinants did not remove the 140,000  $M_r$  band. In view of the strong immunogenicity of the AJ2 determinant, preclearing of AJ2 from cell extracts before immunizations may be advisable when antibodies to less immunogenic components are sought.

#### AJ9, AJ17 and AJ425 Antigenic Systems

Ab AJ9, Ab AJ17 and Ab AJ425 recognized heat-labile determinants on most normal and malignant cells. Differences in serological reactivity with specific cell lines suggest that they identify different antigenic

systems. These antibodies have not precipitated detectable components from [ $^3\text{H}$ ]glucosamine-, [ $^{35}\text{S}$ ]methionine-, or  $^{125}\text{I}$ -labelled SK-MG-1 cells.

## DISCUSSION

This study of human malignant astrocytoma has generated a series of mouse monoclonal antibodies that define nine distinct cell surface antigenic systems. These cell surface determinants have been characterized as restricted or widely distributed based on their distribution on a large panel of cultured tumor cell types. The restricted antigens (AJ225, A010, AJ8, A0122) are preferentially expressed on tumors of neuroectodermal origin. The non-restricted antigens (AJ10, AJ9, AJ17, AJ425, AJ2) are found on virtually all malignant cells. Cell surface determinants present only on tumor cells have not been detected in this analysis; the antigens described here have been identified as normal cell surface components by their expression on at least one normal cell type. The non-restricted antigenic systems are expressed on most normal cell types and are found in brain. The restricted systems, on the other hand, have a restricted distribution on normal cells as well. A010 and A0122 are detected in brain, AJ225 and AJ8 are not. The failure to identify the AJ225 and AJ8 determinants in brain suggests that within the context of normal neural cells AJ225 and AJ8 represent "tumor" antigens. However, end-point absorptions with homogenates of brain may not be sufficiently sensitive to detect antigens restricted to a small subpopulation of neural cells. Extending the serological analysis to sections of normal brain and astrocytomas will be important in assessing the significance of antigens detected on tumor cells in culture but not demonstrable in normal brain by absorption techniques or binding assays.



In this study, subsets of cultured astrocytomas have been identified. The biological significance of grouping astrocytomas on the basis of cell surface characteristics is unknown. Observations in other tumor systems suggest that the surface phenotype of transformed cells reflects both the cell of origin and the state of differentiation of the normal counterpart at the time of malignant change (10). Phenotypic differences among astrocytoma lines may be explained in one of three ways. Astrocytoma subsets may point to transformation in one of several developmentally and phenotypically distinct astrocyte lineages. Alternatively, tumor subsets may indicate that transformation has occurred at different points in a single lineage of astrocyte differentiation. Lastly, tumor subsets may reflect the random loss or random expression of astrocyte differentiation antigens.

The division of cultured astrocytomas into mutually exclusive A010-positive and AJ8-positive subsets and the relationship between the A010/AJ8 surface phenotype and GFA expression provide an opportunity to speculate on the biological significance of these cell surface markers and of the subsets they define. The intriguing feature of A010 and AJ8 expression is the essentially non-overlapping distribution on cultured astrocytomas. It is unlikely that subsetting of this kind is explained by a random expression or loss of the A010 and AJ8 determinants. However, this reciprocal relationship could be taken as evidence for the existence of two antigenically distinct astrocyte lineages or two antigenically distinct phases in a single lineage. Past studies of GFA have indicated that GFA is absent in pluripotential neuroglial stem cells and immature astrocyte precursors but is detectable in all mature astrocytes (11). The observation that A010 is

present on GFA-positive tumors and that AJ8-positive tumors are GFA-negative, suggests that these cell surface markers reflect the state of differentiation of astrocytomas in culture and lends support to the latter view that transformation has occurred in astrocytes at different points in a single developmental lineage.

Serological typing for AJ8, A010 and GFA suggests that cultured astrocytomas can be divided into three groups on the basis of differentiation-related phenotypic characteristics. Cultured astrocytomas which are  $AJ8^-/A010^+/GFA^+$  represent more differentiated cell lines; those which are  $AJ8^+/A010^-/GFA^-$  represent less differentiated cell lines; and those which are  $AJ8^-/A010^+/GFA^-$ ,  $AJ8^-/A010^-/GFA^-$  or  $AJ8^+/A010^+/GFA^-$  represent groups at intermediate stages in differentiation. These relationships are illustrated in Figure 4. It will now be important to determine whether this grouping of cultured astrocytomas on the basis of differentiation characteristics can be shown with tumors in vivo. Antigens AJ225 and A0122 also subset cultured astrocytomas, but the relationship between their expression and other biological properties of normal and malignant astrocytes is uncertain.

The reciprocal expression of A010 and AJ8 also extends to normal cells of neuroectodermal origin. Brain, for example, is  $A010^+/AJ8^-$ ; melanocytes are  $A010^-/AJ8^+$ . Preliminary observations with other neuroectodermal tumors, including neuroblastoma and melanoma, indicate a similar pattern of reciprocal antigen expression. Also noteworthy is the detection of A010 on a T-cell leukemia (MOLT 4), a pattern reminiscent of that observed with other antigens, such as Thy-1, that are shared by T-cells and brain (12).

Because of different serological methods, the use of different cell panels and the limited biochemical characterization of many determinants, especially those that do not precipitate, it is not possible to make direct comparisons between the antigens defined in this report and those described by other investigators in studies of malignant astrocytoma (13). Such comparisons await an exchange of reagents. However, serological differences on a uniform panel of cell lines together with immunochemical differences clearly distinguish the determinants described here from the 12 antigenic systems (gp 150, gp 95, M<sub>19</sub>, R<sub>8</sub>, O<sub>5</sub>, R<sub>24</sub>, gp160, S<sub>25</sub>, gp120r, gp120nr, gp115, V<sub>1</sub>) previously defined in our laboratory by mouse monoclonal antibodies to human malignant melanoma and human renal cancer (1,2,14).

It will be understood that the specification and examples are illustrative but not limitative of the present invention and that other embodiments within the spirit and scope of the invention will suggest themselves to those skilled in the art.

- 17 -  
REFERENCES

1. Dippold, W.G., Lloyd, K.O., Li, L.T.C., Ikeda, H., Oettgen, H.F. and Old, L.J. (1980) Proc. Natl. Acad. Sci. USA 77, 6114-6118.
2. Ueda, R., Ogata, S-I., Morrissey, D.M., Finstad, C.L., Szkudlarek, J., Whitmore, W.F., Oettgen, H.F., Lloyd, K.O. and Old, L.J. (1981) Proc. Natl. Acad. Sci. USA 78, 5122-5126.
3. Pfreundschuh, M., Shiku, H., Takahashi, T., Ueda, R., Ransohoff, J., Oettgen, H.F. and Old, L.J. (1978) Proc. Natl. Acad. Sci. USA 75, 5122-5126.
4. Carey, T.E., Takahashi, T., Resnick, L.A., Oettgen, H.F. and Old, L.J. (1976) Proc. Natl. Acad. Sci. USA 73, 3278-3282.
5. Ueda, R., Shiku, H., Pfreundschuh, M., Takahashi, T., Li, L.T.C., Whitmore, W.F., Oettgen, H.F. and Old, L.J. (1979) J. Exp. Med. 150, 564-579.
6. Brown, J.P., Wright, P.W., Hart, C.E., Woodbury, R.G., Hellstrom, K.E. and Hellstrom, I. (1977) J. Biol. Chem. 255, 4980-4983.
7. Natori, T., Law, L.W. and Appella, E. (1977) Cancer Res. 37, 3406-3413.
8. O'Farrell, P.H. (1975) J. Biol. Chem. 250, 4007-4021.

9. Ogata S., Ueda, R. and Lloyd, K.O. (1981) Proc. Natl. Acad. Sci. USA 78, 770-774.
10. Magrath, I.T. (1981) J. Nat. Cancer Inst. 67, 501-514.
11. Juurlink, B.H.J., Fedoroff, S., Hall, C. and Nathaniel, E.J.H. (1981) J. Comp. Neurol. 200, 375-391.
12. Barclay, A.N., Letarte-Muirhead, M. and Williams, A.F. (1976) Nature 263, 563-567.
13. Schnegg, J.F., Diserens, A.C., Carrel, S., Accolla, R.S. and de Tribolet, N. (1981) Cancer Res. 41, 1209-1213.
14. Pukel, C.S., Lloyd, K.O., Travassos, L.R., Dippold, W.G., Oettgen, H.F. and Old, L.J. (1982) J. Exp. Med. 155, 1133-1147.

TABLE 1. DERIVATION OF MOUSE HYBRIDOMAS PRODUCING MONOCLONAL ANTIBODIES REACTING WITH SURFACE

ANTIGENS OF HUMAN MALIGNANT ASTROCYTOMA CELLS

Astrocytoma cell		No. of positive	No. of clones	
Fusion	line used for	No. of	wells/Total no.	isolated and
No.	immunizations	immunizations	of wells	Antibodies
				Characterized
1	SK-MG-1	6	17/480	4
				AJ2 ( 1)*, AJ8 ( 1)*, AJ9 ( 1)*, AJ17 ( 1)*
2	SK-MG-1	6	10/480	3
				AJ60 ( 1), AJ225 ( $\mu$ )* AJ425 ( 1)*
3	SK-MG-1	8	59/360	1
				AJ10 ( 1)*
4	SK-A02	10	20/360	4
				A010 ( 1)*, A050 ( 1), A092 ( 1), A0122 ( 1)*

\*prototype antibodies (see Table 2)

0104014

TABLE 2. SEROLOGICAL CHARACTERIZATION OF NINE PROTOTYPE MOUSE MONOCLONAL ANTIBODIES DETECTING CELL SURFACE ANTIGENS ON HUMAN MALIGNANT ASTROCYTOMA CELLS

Cells	Ab AJ225		Ab A010		Ab AJ8		Ab A0122		Ab AJ10		Ab AJ9		Ab AJ17		Ab AJ425		Ab AJ2	
	Titer*	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer	Titer
x10 <sup>-3</sup> Abs** x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs x10 <sup>-3</sup> Abs																		
<b>Astrocytomas (GFA)**</b>																		
SK-MG-1	(-) 100	+	-	-	10	+	-	-	100	+	100	+	100	+	100	+	10000	+
SK-MG-2	(+) -	10	+	-	-	-	100	+	100	+	1000	+	10	+	-	+	1000	+
SK-MG-3	(-) 10	+	-	-	-	+	1000	+	100	+	1000	+	100	+	100	+	10000	+
SK-MG-4	(-) 100	+	100	+	-	-	1000	+	10	+	1000	+	10	+	10	+	1000	+
SK-MG-7	(-) 100	+	-	-	-	+	100	+	10	+	1000	+	10	+	1	+	1000	+
SK-MG-9	(-) -	+	100	+	-	-	1	+	1	+	100	+	1	+	1	+	1000	+
SK-MG-10	(-) 10	+	-	-	-	-	-	-	100	+	100	+	1	+	-	+	1000	+
SK-MG-11	(-) 10	+	-	-	10	+	1000	+	1	+	1	+	1	+	1	+	1000	+
SK-MG-12	(-) 10	+	10	+	100	+	-	-	10	+	1000	+	100	+	100	+	1000	+
SK-MG-13	(-) 10	+	-	-	100	+	10	+	-	+	100	+	1	+	100	+	10000	+
SK-M5	(-) -	+	-	-	100	+	1	+	100	+	100	+	10	+	10	+	1000	+
SK-A02	(+) -	+	1000	+	-	-	1000	+	-	+	1	+	1	+	10	+	1000	+
T98	(-) 1	+	-	-	-	+	-	-	10	+	10	+	10	+	100	+	1000	+
U178MG	(-) 100	+	-	-	10	+	-	-	100	+	1000	+	1	+	1	+	1000	+
U251MG	(+) 1	+	1000	+	-	-	-	-	100	+	1000	+	10	+	100	+	1000	+
U373MG	(+) 100	+	1000	+	-	-	-	-	100	+	100	+	100	+	100	+	1000	+
<b>Neuroblastoma</b>																		
SK HMC	-	+	1000	+	-	-	-	-	-	-	1000	+	100	+	100	+	1000	+
SK HSH	-	-	100	+	-	-	-	-	-	-	1000	+	-	-	-	+	10000	+

## Melanoma

SK-MEL-13	-	-	-	-	1000	+	100	+	100	+	100	+	-	+	100	+	1000	+
SK-MEL-28	-	-	-	-	1000	+	100	+	10	+	100	+	100	+	100	+	1000	+
SK-MEL-29	-	-	-	-	-	-	-	-	1000	+	1000	+	100	+	100	+	10000	+
SK-MEL-31	-	-	-	+	-	-	1000	+	100	+	100	+	1	+	10	+	1000	+
SK-MEL-33	-	-	-	+	-	-	-	-	100	+	100	+	100	+	100	+	100	+
SK-MEL-37	-	-	10	+	-	-	100	+	10	+	100	+	10	+	10	+	1000	+
SK-MEL-44	100	+	-	-	-	-	100	+	100	+	10	+	-	+	1	+	100	+
SK-MEL-93	-	-	-	-	1000	+	100	+	1000	+	100	+	1	+	100	+	1000	+
SK-MEL-124	-	-	-	-	-	-	100	+	100	+	100	+	100	+	100	+	100	+
Мелю	-	-	-	-	10000	+	100	+	-	-	1000	+	10	+	10	+	1000	+

## EPITHELIAL CANCERS

21

## Lung

SK-LL-LC	-	-	-	-	-	-	-	-	1	+	1	+	-	+	100	+
SK-LC-6	10	+	-	-	-	-	100	+	10	+	10	+	100	+	10000	+

## Breast

AlAb	-	-	-	-	-	-	-	-	1000	+	10	+	100	+	1000	+
BT-20	-	-	-	-	-	-	-	-	1000	+	10	+	1	+	100	+
CAMA	-	-	-	-	-	-	-	-	-	-	100	+	100	+	1000	+
MCF-7	-	-	-	-	-	-	-	-	100	+	10	+	100	+	1000	+
SK-BR-3	-	-	-	-	-	-	-	-	100	+	10	+	100	+	1000	+

## Colon

HIF-29	-	-	-	-	-	-	-	-	100	+	1	+	-	+	1000	+
SW-1116	-	-	-	-	-	+	-	-	100	+	1	+	-	+	1000	+
SW-1222	-	-	-	-	-	+	-	-	100	+	1	+	-	+	1000	+

0104014



[illegible]

NORMAL HUMAN CELLS											
Brain											
Adult	-	+	-	+	+	+	+	+	+	+	+
Fetal	-	+	-	+	+	+	+	+	+	+	+
Adult skin fibroblasts											
Adult skin	-	-	-	+	100	+	-	+	1	+	10
Fetal skin fibroblasts											
Fetal skin	-	-	-	+	10	+	-	+	100	+	1000
Adult kidney epithelium											
Adult kidney	-	-	-	-	-	100	+	100	+	100	+
Melanocytes											
Melanocytes	1	-	10	100	1000	1000	100	100	10	100	1000
Erythrocytes											
Erythrocytes	-	-	-	-	-	-	-	-	-	-	-

\*Titer: - indicates no reaction in direct tests at a dilution of 1:100.

\*\*Abs: absorption tests. Sera (diluted to end point) were absorbed with the indicated cell type and tested for residual activity for SK-MG-1 (Ab AJ225, Ab AJ10, Ab AJ9, Ab AJ17, Ab AJ425, Ab AJ2), U251MG (Ab A010), or SK-MEL-28 (Ab AJ8, Ab A0122); +, complete absorption; -, no absorption.

\*\*\*GFA: Glial fibrillary acidic protein expression was determined by reactivity of cultured astrocytoma cells (formaldehyde/acetone fixed) with rabbit anti-GFA antiserum (dilution: 1/500) in indirect immunofluorescence tests.

WHAT IS CLAIMED IS

1. Monoclonal antibodies recognizing cell surface antigens on astrocytoma tumor cells.
2. Antibody producing hybridoma cell line characterized by the production of the antibodies as claimed in claim 1.
3. The antibodies of claim 1 wherein the tumor cells are of neuroectodermal origin.
4. Antibody producing hybridoma cell line formed by fusing a myeloma derived cell strain with splenocytes derived from BALB/c mice immunized with human astrocytoma tumor cells.
5. Method of forming an antibody producing hybridoma cell line by fusing a myeloma derived cell strain with splenocytes derived from BALB/c mice immunized with human astrocytoma tumor cells.
6. Method of phenotyping astrocytoma tumor cells comprising determining the reaction of said cells to various phenotyped monoclonal antibodies to astrocytoma tumor cells.

7. A hybridoma for use in the preparation of a monoclonal antibody as claimed in claim 1 and characterised by any one of the ATCC depositions having numbers:-

HB 8338  
HB 8339  
HB 8340  
HB 8341  
HB 8342  
HB 8343  
HB 8344  
HB 8345  
HB 8346  
HB 8347  
HB 8348  
HB 8349